

Assignment 4, STAT 582, due March 24

1. Let (Ω, \mathcal{B}) be a measurable space and let μ and ν be measures. Are the following statements true or false? Give proofs or counterexamples.

- a. If $\mu \ll \nu$, then $\mu(B) \leq \nu(B)$ for all $B \in \mathcal{B}$.
- b. If $\mu \ll \nu$ and $\nu \ll \mu$, then $\mu(B) = \nu(B)$ for all $B \in \mathcal{B}$.
- c. If $\nu \ll \mu$ and μ is not σ -finite, then $\frac{d\nu}{d\mu}$ does not exist.
- d. $\frac{d\nu}{d\nu}$ always exists and equals 1.
- e. If $\nu \ll \mu$ and μ is σ -finite, then ν is σ -finite.

2. Let μ be counting measure on $(\mathbb{R}, \text{Borel})$.

a. Let λ denote Lebesgue measure. Show that $\lambda \ll \mu$ but that $d\lambda/d\mu$ does not exist. Why does this not contradict the Radon-Nikodym theorem?

b. Describe the class of measures on $(\mathbb{R}, \text{Borel})$ that are absolutely continuous with respect to μ .

c. Is $\mu \ll \lambda$?

3. Let (Ω, \mathcal{B}, P) be a probability space and let $\mathcal{G} \subseteq \mathcal{B}$ be a sub- σ -field of \mathcal{B} . Let A, A_1, A_2, \dots be disjoint sets in \mathcal{B} . Show that

- a. $P(A^c | \mathcal{G}) = 1 - P(A | \mathcal{G})$
- b. $P(\bigcup_{n=1}^{\infty} A_n | \mathcal{G}) = \sum_{n=1}^{\infty} P(A_n | \mathcal{G})$.

4. Let $\Omega = \{-1, 1\}$, $\mathcal{B} = \{\emptyset, \{-1\}, \{1\}, \Omega\}$ and P such that $P(\{-1\}) =$

$P(\{1\}) = 1/2$. Further, let $X(\omega) = \omega$, $Y(\omega) = 2\omega$, $g(x) = x^2$ and consider $\mathcal{G}_1 = \sigma(Y)$ and $\mathcal{G}_2 = \sigma(g(Y))$.

a. What are \mathcal{G}_1 and \mathcal{G}_2 ?

b. Find $E[X|\mathcal{G}_1]$, $E[X|\mathcal{G}_2]$, $E[E[X|\mathcal{G}_1]|\mathcal{G}_2]$ and $E[E[X|\mathcal{G}_2]|\mathcal{G}_1]$. Verify the "smallest σ -field always wins" property.

5 Let $X \sim \text{unif}[-1, 2]$. Show that $E[X|X^2] = X \cdot I_{\{X>1\}}$.