

# STAT 552 Homework 3

Due date: In class on Tuesday, October 3, 2006

Instructor: Dr. Rudolf Riedi

9. A Markov chain has state space  $S = \{1, 2, \dots, 8\}$ . Starting from  $X_0 = 1$ , the chain moves in each step from its current state  $j$  to any of the larger states  $\{k : k > j\}$  with equal probability. State 8 is absorbing.

- (a) Compute the transition matrix.
- (b) Decompose the state space into recurrent and transient classes.
- (c) Find the expected number of steps to reach state 8.

Hint: Set  $N = \inf\{n \geq 0 : X_n = 8\}$  and  $q_i = \mathbb{E}[N|X_0 = i]$  and show first the following:

$$q_i = \mathbb{E}[N|X_0 = i] = \sum_{k=1}^8 \mathbb{E}[N|X_0 = i, X_1 = k]p_{ik} = \sum_{k=1}^8 (1 + q_k)p_{ik}. \quad (1)$$

Here you need to justify both, steps 2 and 3. Finish by solving the recursion starting at  $q_8$  which is easy (trivial) to compute. You should find  $q_1 = 363/140 = 2.6$  (pocket calculator may be handy but not required).

10. Let  $X_n$  be a Markov Chain.

- (a) Assume that  $f$  is a given 1-1 function of the state space, i.e.,  $f$  is invertible. Show that the sequence of random variables  $f(X_n)$  form a Markov Chain as well.
- (b) Show that if  $f$  is not invertible then this can be true or false, depending on the chain and the function. Hint: Consider  $f(x) = x^2$  or  $f(x) = |x|$  and a small state space. Give an example of a MC  $X_n$  where  $f(X_n)$  is not Markov, and one where  $f(X_n)$  is Markov.

11. Let  $\mathbf{P} = [p_{ij}]_{(ij)}$  denote the transition matrix of an MC. Define  $q_{ij} = 1$  if  $p_{ij} \neq 0$  and  $q_{ij} = 0$  else. The matrix  $\mathbf{Q} = [q_{ij}]_{(ij)}$  indicates whether it is possible to reach  $j$  from  $i$  in 1 step.

- (a) Show that the matrix  $\mathbf{Q}^2$  indicates in how many ways it is possible to reach  $j$  from  $i$  in 2 steps.
- (b) Assume that  $S$  has  $m$  states. Explain how the matrix  $\mathbf{Q} + \mathbf{Q}^2 + \dots + \mathbf{Q}^m$  can be used to decide whether  $j$  is reachable from  $i$  or not.  
Hint: Using a recursive argument show that  $\mathbf{Q}^k$  indicates in how many ways it is possible to reach  $j$  from  $i$  in  $k$  steps.

12. (a) Assume that there exists an integer  $n$  such that  $p_{ij}^{(n)} \neq 0$  for all  $i, j \in S$ . Show that the MC is then irreducible (meaning that  $S$  forms class).

- (b) \*Bonus question\* The reverse is not true. Give a simple counter example.