

STAT 582 Homework 1

Due date: In class on Friday, February 4, 2005

Instructor: Dr. Rudolf Riedi

1. Let X_n be a monotone sequence of random variables. Assume that $X_n \xrightarrow{P} X$. Show that $X_n \xrightarrow{\text{a.s.}} X$.
Hint: Consider subsequences.

2. Let X_n be any sequence of random variables. Show that

$$X_n \xrightarrow{\text{a.s.}} X \iff \sup_{k \geq n} |X_k - X| \xrightarrow{P} 0.$$

Hint: this is very similar to a result developed in class.

3. Give an example of a probability space such that for all $0 < p < 1$ the expression $\|X\|_p = (\mathbb{E}|X|^p)^{1/p}$ is not a norm, i.e., such that the triangular inequality $\|X + Y\|_p \leq \|X\|_p + \|Y\|_p$ fails for some random variables X and Y .

Hint: Consider a simple coin toss, and set X and Y such that $X + Y = 1$ always.

4. Let X_n be any sequence of random variables. Set $S_n = X_1 + \dots + X_n$.

(a) Assume that the sequence of real numbers a_n converges to a . Show that then also $(a_1 + \dots + a_n)/n$ converges to a .

(b) Conclude that if $X_n \xrightarrow{\text{a.s.}} 0$ then $S_n/n \xrightarrow{\text{a.s.}} 0$.

(c) Let $p \geq 1$. Show that if $X_n \xrightarrow{L_p} 0$ then $S_n/n \xrightarrow{L_p} 0$. Hint: triangular inequality.

(d) Show that $X_n \xrightarrow{P} 0$ does NOT imply $S_n/n \xrightarrow{P} 0$.

Hint: Consider a suitable sequence of random variables with $X_n = 2^n$ with probability $1/n$, and $X_n = 0$ else.