

**1. [20 points]** For each of the statements below, circle **T** or **F** for “True” or “False,” respectively. (4 pts. each)

**T F** : The correlation can be any number, but it is usually between  $-1$  and  $+1$ .

**FALSE: Correlation must be between  $-1$  and  $+1$ .**

**See item 1. at the bottom of p. 166 in the text.**

**T F** : If the distribution is bell shaped with no outliers, we expect *IQR* will be smaller than  $s$ .

**FALSE: It was stated in class that *IQR* is approximately equal to  $1.35s$  for the normal distribution. You can also figure this out from the tables.**

**T F** : Nonresponse bias refers to systematic error in sampling from a population due to subjects being unavailable or refusing to reply.

**TRUE: See bottom of p. 251 in the text.**

**T F** : One purpose of randomization in experimental design is to eliminate confounding effects from lurking variables that might be present in an observational study.

**TRUE: This was stated in lecture.**


**T F** : Both the mean and the median are resistant measures of the center of a distribution of data.

**FALSE: See the bottom of p. 37 to the top of p. 38.**

**2. [30 points]** Use the table below to sketch a density histogram for the data.

Class	Percentage		Class Width	Bar Height
0 – 20	10%	→	<b>20</b>	<b>0.5</b>
20 – 30	20%	→	<b>10</b>	<b>2.0</b>
30 – 40	30%	→	<b>10</b>	<b>3.0</b>
40 – 60	20%	→	<b>20</b>	<b>1.0</b>
60 – 100	20%	→	<b>40</b>	<b>0.5</b>

We have added two extra columns in the table : one for class width and one for the height of the histogram bars. The plot appears below.



**3. [30 points]** Suppose a data set has approximately a normal distribution with mean  $\bar{x} = 200$  and standard deviation  $s = 20$ .

(a) Estimate the percentage of the data which are between 170 and 210.

**Computing the corresponding  $z$  values:**


$$z_1 = \frac{170 - 200}{20} = -1.5 \quad (1)$$

$$z_2 = \frac{210 - 200}{20} = 0.5 \quad (2)$$

From the tables provided, the area under the curve to the left of  $z_1 = -1.5$  is 0.0668, and the area under the curve to the left of  $z_2 = 0.5$  is 0.6915. Thus, the area between them is

$$0.6915 - 0.0668 = 0.6247.$$

The calculation is depicted pictorially below.



**3(b)** Find approximately the 80'th percentile of the data.

Using the tables, the 80'th percentile of the  $N(0,1)$  distribution is 0.84. The area corresponding the  $z = 0.84$  is 0.7995, which is the closest we can get to 0.8. The corresponding data value is obtained by the “inverse”  $z$ -value transformation:

$$x = \bar{x} + z * s = 200 + 0.84 * 20 = 216.8.$$

Thus, our estimate of the 80'th percentile of the data is 216.8.

**4. [20 points]** Below are 5 values of  $r$ , the correlation of a sample, and 4 scatterplots. Match the value of  $r$  with the scatterplot by writing the plot label (A, B, C, or D) next to the value of  $r$ . Obviously, one value of  $r$  will be unmatched.

(i)  $r = -0.9$  Plot C. Clearly this plot has a negative association and the points fall close to a straight line, so the corresponding correlation is near  $-1$


(ii)  $r = -0.5$  No Plot.

(iii)  $r = 0.0$  Plot B. There is no upward or downward “tilt” in Plot B, so there is no linear association, although there is a clear nonlinear association.


(iv)  $r = +0.5$  Plot A. Plot A displays a fairly clear upward tilt, hence has a positive correlation, but it is not so strongly positive as Plot D, so by process of elimination Plot A must go with  $r = +0.5$ .

(v)  $r = +0.9$  Plot D. Plot D has a very strong positive correlation, near 1.


Plot A




Plot B



Plot C



Plot D



## Summary Statistics for Scores

For  $n = 75$  persons taking the exam before Fri., 13 Feb.:

$$\bar{x} = 87.01, \quad s = 13.07.$$

Five Number summary:

min	$Q_1$	$M$	$Q_3$	max
56	77	92	99	100

Histogram is shown below. Approximate letter grades are

A-B	85 – –100	(68%)
C	65 – 84	(25%)
D	56 – 64	(7%)

Histogram of Exam1 Scores

